arXiv:1209.4433v1 [math.OC] 20 Sep 2012

Contraction Criteria for Existence, Stability, and Robustness of a Limit Cycle

Ian R. Manchester[1] Jean-Jacques E. Slotine[2]

1: School of Aerospace, Mechanical and Mechatronic Engineering, University of Sydney, Australia
2: Nonlinear Systems Laboratory, Massachusetts Institute of Technology, USA
ian.manchester@sydney.edu.au jjs@mit.edu

Abstract— **Stable periodic behavior is the desired behavior of many robotic systems, such as walking robots, swimming robots, and other biologically inspired systems. Designing and verifying controllers for stable periodic behaviour is significantly more challenging than for equilibria. We study pointwise contraction conditions for an autonomous system which imply the existence and global stability of a limit cycle in a compact manifold. Unlike methods based on Lyapunov theory, our condition does not require knowledge of the location of the limit cycle in state-space. This allows straightforward extension to robustness via parameter-dependent certificates, and compositions of systems in feedback and serial interconnections. This new criterion also has interesting connections to contraction theory for trajectories, orbital stability analysis via transverse linearization, and topological facts about regions of attraction of limit cycles.**

I. INTRODUCTION

Systems with periodic solutions are central to biologically-inspired robot locomotion [1], [2], and are also in many other areas of science and engineering, e.g. phase-locked loops or vortex shedding from aircraft. In biology, oscillating systems seem to be the rule rather than the exception [3].

The basic question we address in this paper is the following: when does a system of the form

$$\dot{x} = f(x) \tag{1}$$

have the property that all solutions starting from a particular set K converge asymptotically to a unique limit cycle? This is the property of orbital stability, and is substantially more difficult to deal with than simple stability of an equilibrium.

There is a long and distinguished history of research into limit cycles for nonlinear systems. For example, the famous result of Poincaré-Bendixson gives a very simple condition for planar systems. However this result relies heavily on the Jordan curve theorem and for higher-dimensional systems fails to hold: bounded orbits of smooth systems may have chaotic behaviour. An important generalization to monotone cyclic feedback systems was published in [4], however this depends on quite a special system structure and there are many application areas where it does not apply. Birkhoff gave necessary conditions for periodic solutions in terms of the existence of particular "phase variables", or associated differential one-forms [5], [6].

However, all of these conditions imply the existence of *at least one* limit cycle, but give no insight into the number of limit cycles, or their stability. Previous work by the first author and colleagues has developed the Lyapunov-based "transverse dynamics" approach to study regions of stability of limit cycles [7], [8], [9], [10], however this approach is not applicable when the system dynamics are uncertain, since uncertainty will generally change the location of the limit cycle in state space.

In recent years many efficient computational methods for proving stability of equilibria of nonlinear systems have been proposed, based on searching for "stability certificates" such as Lyapunov functions and contraction metrics. For the purposes of robust stability analysis, a central difference between these two approaches is that a Lyapunov function must generally be constructed about a known equilibrium, whereas a contraction metric implies the existence of an equilibrium indirectly [11]. This can be useful for robustness analysis of highly nonlinear systems where the equilibrium point may change depending on the unknown dynamics [12].

In this paper we extend contraction analysis to the study of limit cycles. It will be shown that the existence of a contraction metric of a particular "transverse" form is equivalent to the existence of a unique orbitally stable limit cycle. In other words, we decouple the property of *convergence* from the task of finding a particular limiting solution. A major benefit of this approach is that conditions for stability can be extended to conditions for *robustness*, i.e. existence of a stable limit cycle under parameter uncertainty and interconnection with other (possibly uncertain) systems.

II. PROBLEM SETUP AND PRELIMINARIES

We assume that $f : K \to \mathbb{R}^n$ in (1) is smooth and $x \in \mathbb{R}^n$. In the situations we will consider in this paper the solution of (1) will always exist and be unique.

A periodic solution $x^\star$ is one for which there exists some $T > 0$ such that $x^\star(t) = x^\star(t + T)$ for all t. Equilibria are trivially periodic for every T, but for oscillatory solutions – which are our main concern – there is some minimal time T such that the above holds and this is referred to as the *period*. The *orbit* of a periodic solution is the set $\mathcal{X}^\star := \{x : x = x^\star(t) \text{ for some } t\}$. Note that while periodic *solutions* cannot be asymptotically stable, their *orbits* can be, and in this case we may say that the solution is *orbitally stable* (see, e.g.,



Fig. 1. Regions of attraction to limit cycles are homeomorphic to discs parametrized by a scalar phase variable τ.

[13]). We will also refer to this as "convergence to a periodic orbit".

The basic statement is: give pointwise conditions on the vector field $f(\cdot)$ and a compact set $K\subset\mathbb{R}^n$ guaranteeing that every solution of (1) starting from an initial condition in K converges to a unique periodic orbit $\mathcal{X}^\star$ as $t\to\infty$.

III. STABILITY AND REGIONS OF ATTRACTION OF PERIODIC SOLUTIONS

In this section we discuss some aspects of oscillating systems that render the study of stability substantially more challenging that stationary systems.

It is well known that periodic solutions of an autonomous differential equation can never be locally asymptotically stable. This is clear from the fact that solutions which have initial conditions on the periodic orbit but offset in time will never approach each other. Orbital stability is the strongest form of stability that can be hoped for.

There are also interesting properties of the "global" structure of regions of attraction to periodic orbits. It is known that the the region of attraction is a continuous deformation of the set $\mathbb{D}^{n-1}\times\mathbb{S}^1$, i.e. the cartesian product of an open unit disc of dimension one less than the full state space, with a single circle coordinate. These are often referred to as "transversal" and "phase" coordinates, respectively. In all cases except possibly with $n=5$ it is guaranteed that the deformation is smooth differentiable, due to the recent resolution of the Poincaré Hypothesis [6].

In that case, let us consider such a change of variables:

$$x_\perp=p(x),\quad \tau=q(x) \tag{2}$$

with $p(x)\in\mathbb{R}^{n-1}$ and $\tau\in\mathbb{S}^1$, which we assume to be differentiable. For τ to be considered a "phase variable" it must satisfy $\dot\tau=\frac{dq}{dx}f(x)>0$ for all $x\in K$.

Indeed, the existence of a global phase variable – a scalar variable that is always increasing – implies the existence of at least one periodic orbit, though not stability [5].

A change of variables into phase and transversal coordinates depends on the existence of a *codimension-1 foliation*, i.e. a family of $n-1$-dimensional submanifolds ("leaves") which are parametrized by the phase variable. Not all nonsingular vector fields admit such a foliation, e.g. Hopf bifurcations do not [14]. Regardless, even if such a transformation exists it may be prohibitively hard to *find*.

Nevertheless, let us mention briefly some important points. From the definition of the transformations, we have

$$\dot x_\perp=\frac{\partial p}{\partial x}(x)f(x),$$

and

$$\dot\tau=\frac{\partial q}{\partial x}(x)f(x),$$

noting that $\dot\tau$ is a scalar greater than zero.

It is clear that τ can be considered a reparametrization of time, and then one can analyze $\frac{dx_\perp}{d\tau}$ instead of $\dot x_\perp$. Since $\tau\in\mathbb{S}^1$ the differential equation

$$\frac{dx_\perp}{d\tau}=\frac{1}{\frac{\partial q}{\partial x}(x)f(x)}\frac{\partial p}{\partial x}(x)f(x)=:f_\perp(x_\perp,\tau)$$

can be considered to be a periodically time-varying differential equation of dimension $n-1$. Now, it is known that systems which are periodically time-varying and *contracting* have the property that all solutions converge to a particular periodic solution [11]. Making the "lifting" transformation back to considering τ to be a state, this then implies convergence to a periodic *orbit* in dimension n. Thus if one can construct a transformation to phase and transversal coordinates, and prove that the transversal system is contracting, then one has proven orbital stability.

The remarkable fact is that one can show *transverse contraction* without ever constructing the global change of variables.

IV. CONTRACTION CONDITIONS FOR LIMIT CYCLES

The following theorem relates existence of a special form of metric to a property sometimes referred to as Zhukovsky stability, which is deeply linked to the existence of periodic solutions [15].

Theorem 1: Let $K\subset\mathbb{R}^n$ be compact, path-connected, and strictly forward invariant. If there exists a smooth positive-definite matrix function $M(x)$ on K and a $\lambda>0$ such that

$$\delta'(\dot M+MF+F'M+\lambda M)\delta\leq 0, \tag{3}$$

for all $\delta\in\mathbb{R}^n$ satisfying $\delta'M(x)f(x)=0$, then for every two solutions x_1 and x_2 with initial conditions in K there exists time reparametrizations $\tau_1(t)$, $\tau_2(t)$ such that $x_1(\tau_1(t))\to x_2(\tau_2(t))$ as $t\to\infty$.

Proof: since the set K is path-connected by definition there exists a path between any two points x_0 and x_1 in K. Such a path can be considered as a smooth mapping $\gamma:[0,1]\to\mathbb{R}^n$ with $\gamma(0)=x_0$ and $\gamma(1)=x_1$.

Denote by $\Gamma(x_0,x_1)$ the set of all such smooth paths between x_0 and x_1 remaining in K and associate with each a length

$$L(\gamma)=\int_0^1\left(\frac{\partial}{\partial s}\gamma(s)\right)'M(\gamma(s))\frac{\partial}{\partial s}\gamma(s)ds$$

and then the standard Riemannian metric between two points x_0 and x_1 is

$$d(x_0, x_1) = \inf_{\gamma \in \Gamma(x_0, x_1)} L(\gamma)$$

As in contraction analysis, we study the instantaneous change in the length $L(\gamma)$ as x_1 and x_2 evolve along solutions of $\dot{x} = f(x)$:

$$\frac{d}{dt}\left(\frac{\partial}{\partial s}\gamma(s)\right)' M(\gamma(s))\frac{\partial}{\partial s}\gamma(s)$$

However, since we are allowing time reparametrizations between solutions, we also allow each point on $\gamma(s)$ to move at a different rate $\frac{d\gamma(s)}{dt} = f(\gamma(s))a(\gamma(s))$. Then

$$\frac{d}{dt}\frac{\partial \gamma}{\partial s} = F(\gamma(s))a(s) + f(\gamma(s))\frac{\partial a(\gamma(s))}{\partial s}$$

Without loss of generality for this pointwise condition we can let $a(s) = 1$ and denote $z := \frac{\partial a(\gamma(s))}{\partial s}$, then in terms of the virtual displacement δ we have

$$\dot{\delta} = F(x)\delta + zf(x)$$

$$\begin{aligned}\frac{d}{dt}(\delta' M\delta) =& \delta'\left(F(x)'M(x) + M(x)F(x) + \dot{M}(x)\right)\delta \\ &+ 2z\delta' M(x)f(x)\end{aligned}$$

Now, since z is a free variable, there always exists a z making the evolution of this length segment negatative, except possibly when $\delta' M(x)f(x) = 0$ and z has no influence. From this we derive the transverse contraction condition in the theorem. □

It is known that systems satisfying such a property have limit cycles [15], but with the framework of contraction the proofs are simpler, so we give a proof here.

Theorem 2: If the conditions of Theorem 1 are satisfied, then all solutions starting with $x(0) \in K$ converge to a unique limit cycle.

Proof: since K is invariant and compact, it follows that $\Omega(x_0)$ exists and is a compact subset of K. Furthermore, a clear implication of Theorem 1 is that all points in K have the same ω-limit set, which we denote $\Omega(K)$.

Pick a point $x^\star$ in $\Omega(K)$. By strict invariance, this is an interior point of K. By assumption, $f(x^\star) \neq 0$ so we can construct the hyperplane orthogonal to $f(x^\star)$, which we denote S. We will prove convergence to a limit cycle by constructing a Poincaré map on S.

Since $f(\cdot)$ is smooth, for x in some neighborhood B of $x^\star$ we have that $f(x)'f(x^\star) > 0$, so in $B_S := B \cap S$ solution curves are transversal to S and pass through it in the same direction as at $x^\star$.

Since $x^\star$ is in the ω-limit set for all points in K, and B_S is transversal, the evolution of the system from any point $x_0(t) \in B_S$ eventually passes through B_S again, i.e. $x_0(t+s) \in B_S$ where $s > 0$ depends on x_0. This evolution can be represented by a Poincaré map $T : B_S \to B_S$.

Take the distance between two points $d(x_0, x_1)$ on B_S to be Riemannian metric distance from Lemma 1. Note that although the two points lie on the $n-1$ dimensional set B_S, the curves joining them in the definition of d may pass out of the plane and through n-dimensional space. By Lemma 1, we have that $d(T(x_0), T(x_1)) < d(x_0, x_1)$. Hence T is a contractive map from B_S onto itself, and by the Banach fixed point theorem has a unique stable fixed point, which is its only limit point so must be $x^\star$. By standard results on Poincaré maps this implies that $x^\star$ is a point on a limit cycle, which by Proposition 1 all solutions converge to.
□

The new conditions is weaker than contraction, which implies that any two solutions converge, i.e. $x_0(t) \to x_1(t)$, but stronger than *orbital stability*, which implies only convergence to the same ω-limit set and thus doesn't preclude chaos [15]. Our condition implies a behaviour that is sometimes referred to as Zhukovsky stability and has been used in several recent papers on limit cycle stability and system identification, see e.g. [16], [9], [17] and apparently goes back to Poincaré in its essential argument [13].

V. SINGULAR CONTRACTION METRICS

The following equivalent condition is easier to use in some applications, where a set of transversal coordinates are available [9].

Theorem 3: Given a particular choice of normal vector $z(x)$ satisfying $z(x)'f(x) > 0$ everywhere[1] and a particular set of transversal coordinates, condition (3) is equivalent to the existence of a singular positive-semidefinite contraction metric $M_s(x)$ with rank$[M_s(x)] = n-1$ and transversal, i.e. $M_s(x)z(x) = 0$, which satisfies

$$\dot{M}_s + M_sF + F'M_s + \lambda M_s \leq 0. \tag{4}$$

Proof: Let $\pi(x) \in \mathbb{R}^{1\times n}$ be the orthogonal projection matrix on to the subspace spanned by $z(x)$. Let $\Pi(x) \in \mathbb{R}^{(n-1)\times n}$ be the projection on to the subspace orthogonal to $z(x)$, i.e. on to the transversal coordinates. Let $f_\perp = \Pi(x)f(x)$ and $f_\| = \pi(x)f(x)$ be the projections of $f(x)$ on to these subspaces.

Decompose the singular metric like so:

$$M_s(x) = \Pi(x)'\bar{M}(x)\Pi(x),$$

then we can construct a new positive-definite metric $M(x)$ in the following way

$$M(x) = \begin{bmatrix}\Pi(x) \\ \pi(x)\end{bmatrix}' \begin{bmatrix}\bar{M}(x) & -\bar{M}f_\perp/f_\| \\ -f_\perp'\bar{M}/f_\| & 1\end{bmatrix}\begin{bmatrix}\Pi(x) \\ \pi(x)\end{bmatrix}.$$

The interpretation of this is that a differential δ is decomposed into a component collinear with $f(x)$ which we can denote by $\delta_\|$ and a component orthogonal to this denoted by $\delta_\perp$. Then

$$\delta' M(x)\delta = \delta_\perp'\bar{M}(x)\delta_\perp - 2\delta_\perp'\bar{M}f_\perp\delta_\|/f_\| + |\delta_\||^2.$$

Note that $f_\| \neq 0$ since $f(x) \neq 0$ and $f(x)'z(x) > 0$. Now, the conditions of Theorem 1 are that $\delta' M(x)f(x) = 0$.

---

[1] Taking $z(x) = f(x)/|f(x)|$ is one possible choice, but in some applications freedom in $z(x)$ is beneficial.

By the above construction of $M(x)$, the terms involving $\delta_\perp$ cancel and so $\delta' M(x) f(x) = 0$ implies

$$\delta_{\|}(-f_\perp' \bar{M} f_\perp + f_{\|}^2) = 0$$

But since M_s is positive definite, and $f_\perp$ and $f_{\|}$ are projections on to an orthonormal basis of $f(x)$, and $f(x) \neq 0$, it follows that the term in the brackets can not be zero, so this implies that $\delta_{\|} = 0$. This implies that

$$\frac{d}{dt}\left(\delta' M(x) \delta\right) = \frac{d}{dt}\left(\delta_\perp' M_s(x) \delta_\perp\right)$$

which by Condition (4) is negative, and hence the system satisfies the conditions of Theorem 1.

Given a metric $M(x)$ satisfying the conditions of Theorem 1, there exists a stable periodic solution which all initial conditions converge to. By constructing a differential system explicitly in the transversal coordinates defined be the condition $z(x)'\delta = 0$ using the method described in [9] one has a stable time-varying linear system of dimension $n-1$. By standard results on Lyapunov functions for time-varying linear systems, along each time-reversed trajectory there exists a smooth solution to the Lyapunov differential equation, which can be used as a singular contraction metric. See [9] for more details on such constructions.

□

This formulation is also easy to extend to hybrid dynamical systems (e.g. walking robots) by enforcing a discrete contraction condition on switching surfaces, see [11] and[9] for details.

A. Generalized Jacobian and Transverse Linearization

In this section we show given the existence of a singular contraction metric, the negative definiteness of an associated generalized Jacobian is equivalent to contraction of transversal coordinates.

Let

$$M_s(x) = \Pi(x)' \bar{M}(x) \Pi(x)$$

where $\bar{M}(x) \in \mathbb{R}^{n-1}$ and is positive definite. $\Pi(x) : \mathbb{R}^n \to \mathbb{R}^{n-1}$ is a projection on to the subspace orthogonal to $f(x)$.

Consider a decomposition of $\bar{M}(x) = L(x)' L(x)$ with $L(x)$ nonsingular, then set $\Theta(x) := L(x)\Pi(x)$ and consider the generalized Jacobian of the system under the change of coordinates $z = L(x)\Pi(x)x$

$$\dot{\delta}_z = \bar{F} = (\dot{\Theta} + \Theta F)\Theta^\sharp \delta_z$$

where $\sharp$ denotes pseudoinverse. Note that $\Theta^\sharp = \Pi(x)' L(x)^{-1}$. Then we have

$$\bar{F} = \dot{L}\Pi\Pi' L^{-1} + L(\dot{\Pi}\Pi' + \Pi F \Pi')L^{-1}.$$

The term $\dot{\Pi}\Pi' + \Pi F \Pi'$ is exactly the transverse linearization from [9] with orthogonal hypersurfaces, which we denote $F_\perp$. In fact, in the trivial case in which the identity is a contraction metric, the generalized Jacobian is the transverse linearization. Now, since Π is a projection on to a lower dimensional space, $\Pi\Pi' = I$, so the above can be simplified as

$$\bar{F} = \dot{L}L^{-1} + L F_\perp L^{-1}$$

If the generalized Jacobian is uniformly negative:

$$\bar{F} + \bar{F} \leq -\lambda I$$

for some $\lambda > 0$, we can multiply both sides of the above inequality by L and obtain

$$\dot{L}'L + L'\dot{L} + L'LF_\perp + F_\perp' L'L \leq -\lambda L'L$$

And since $\bar{M} = L'L$ this is

$$\dot{\bar{M}} + \bar{M}F_\perp + F_\perp'\bar{M} \leq \lambda \bar{M}$$

which is exactly the conditions for contraction of the transverse dynamics [11]. Furthermore, when a periodic solution is known, then this is also a Lyapunov ODE condition, which is satisfied if and only if the periodic solution is orbitally exponentially stable [9].

Note that in [9] the transverse dynamics were computed explicitly about a known periodic solution and used to compute regions of orbital stability. In this paper, we consider the linearization of the transverse dynamics about *every* solution of the system, though only implicitly. In general, it is difficult to compute an analytical formula for $L(x)$: in principle it could be done via the Cholesky algorithm although the expressions would be extremely complicated.

Given the results of [11], the following theorem is an easy consequence of the above derivation:

Theorem 4: If there exists a $\Pi(x)$, L so that $\bar{F}$ is contracting with respect to some matrix measure on a compact forward-invariant set, then there exists a unique orbitally stable limit cycle.

VI. EXTENSIONS OF THE BASIC THEORY

The results in the previous sections are derived for autonomous systems, i.e. time-invariant systems with no exogenous inputs. However, the transverse contraction machinery can be applied for a much wider range of systems including compositions, feedback interconnections, and systems with bounded disturbances.

In many applications in which exact model parameters are unavailable, it is desirable to characterize parameter ranges over which the qualitative behaviour of the system remains the same. Engineering motivations are well known, but robustness analysis has also become of interest recently in biology, including as a measure of model validity [23]. In particular, the papers [23], [24] address robustness of limit cycles by gridding over parameter ranges and simulating the nonlinear system until convergence can be ascertained. Gridding and simulation becomes very expensive computationally as the number of parameters grows, so alternative methods are desirable.

Robustness analysis of equilibria and trajectories is classical topic for which many methods have been derived, but robustness of limit cycles is substantially more challenging to address. In [25] small-gain conditions were derived for limit-cycle robustness, though such conditions are often very conservative.

A. Feedback Interconnections and Incremental Dissipativity

For this result we consider systems with exogenous inputs:

$$\dot{x} = f(x, w) \tag{5}$$

We define a system as *transverse incrementally dissipative* (TID) with a supply rate $\sigma(x, w, \delta_x, \delta_w)$ if the following condition holds

$$\frac{d}{dt}\left(\delta_x' M(x)\delta_x)\right) \leq \sigma(x, w, \delta_x, \delta_w).$$

for all δ_x such that $\delta_x' M(x) f(x, w) = 0$. Note that if $w = 0$ then the above condition reduces to non-strict transverse contraction. Hence, every TID system is transverse contracting.

If the system (5) also has an output $y = g(x, w)$ with Jacobian $\frac{\partial g}{\partial x} = G(x, w)$, then we can mention two particularly useful informative supply rates: incremental versions of classical small-gain:

$$\sigma_\gamma(\delta_x, \delta_w) = \gamma|\delta_w|^2 - |G(x, w)\delta_x|^2,$$

and passivity:

$$\sigma_p(\delta_x, \delta_w) = \delta_w' G(x, w)\delta_x,$$

where passivity requires that input w and output y have the same dimension.

We now give the following theorem:

Theorem 5: Given two systems $\dot{x}_1 = f_1(x_1, w_1)$ and $\dot{x}_2 = f_2(x_2, w_2)$, which are interconnected so that $w_1 = g_2(x_2)$ and $w_2 = g_1(x_1)$. Then if the two systems are TID with supply rates σ_1, σ_2 such that $\sigma_1 + \sigma_2 \leq -\alpha|\delta_x|^2$ for some $\alpha > 0$, then the interconnected system is transverse contracting.

Proof Let δ_1 and δ_2 refer to the state differentials of system 1 and system 2, respectively, and $\delta_x = [\delta_1 \delta_2]'$. Furthermore, let $f(x) = [f_1(x_1, g_2(x_2))\ f_2(x_2, g_1(x_1))]'$ be the dynamics of the interconnected system.

Suppose the two systems are verified to be TID with metrics $M_1(x_1)$ and $M_2(x_2)$, then consider the metric for the interconnected system

$$M = \begin{bmatrix} M_1(x_1) & 0 \\ 0 & M_2(x_2) \end{bmatrix}$$

It is clear from construction that this satisfies

$$\frac{d}{dt}\left(\delta_x' M \delta_x\right) < -\alpha|\delta_x|^2 \leq \lambda \delta_x' M(x)\delta_x$$

for all δ_x satisfying $\delta_x' M(x) f(x) = 0$ and for some $\lambda > 0$, where the second inequality follows from the compactness of K and continuity and positive-definiteness of $M(x)$. $\square$.

The above method is related to the so-called *S-Procedure* in robust control, widely used in H^∞ and Integral Quadratic Constraint analysis [18], [19].

Note that TID is a stronger property than incremental dissipativity so, for example, any linear passive system satisfies the TID property with supply rate σ_p, and any linear stable system satisfies TID with supply rate σ_γ where γ is the square of the H^∞ norm of the system (peak frequency response). The interconnections of oscillators with passive systems has been studied in the past via describing function methods [20], but without rigorous proofs of stability.

This formulation may also be helpful in analyzing interconnections of oscillators [21], [22]. Note that we have a condition by which two *non-identical* oscillators will, when interconnected, converge to a periodic solution which may be entirely different than their natural oscillations.

B. Serial Compositions of Systems

A relatively simple application of the above theorem is to consider the composition of a contracting system and a transverse-contracting system.

$$\dot{x}_1 = f_1(x_1) \tag{6}$$

$$\dot{x}_2 = f_2(x_1, x_2) \tag{7}$$

Theorem 6: Suppose for each fixed x_1, f_2 is transverse contracting with metric $M_2(x_2, x_1)$, i.e.

$$\delta_2'(F_2' M_{22} + M_2 F_{22} + \frac{\partial}{\partial x_2} M_2 f(x_2) + \lambda_2 M_2)\delta_2 \leq 0$$

for all δ_2 satisfying $\delta_2' M_2 f_2 = 0$ and f_1 is "fully" contracting in the sense of [11], i.e. there exists $M_1(x_1)$ such that

$$F_1' M_1 + M_1 F_1 + \frac{\partial}{\partial x_1} M_1 f(x_1) + \lambda_1 M_1 \leq 0$$

then the composed system is transverse contracting, and hence has a unique stable limit cycle.

Proof we prove this theorem by constructing a metric which decomposes as

$$\delta' M_c \delta := \delta_1' M_1 \delta_1 + \alpha \delta_2' M_2 \delta_2 = 0$$

which will be shown to verify the existence of a unique stable limit cycle. Let $x = [x_1' x_2']'$ and $f(x) = [f_1(x_1)' f_2(x_1, x_2)']'$. Since the contraction conditions are homogeneous with respect to δ, and $\delta = 0$ is trivial, it is sufficient to consider the case where $|\delta| = 1$.

First, we note that since $f_2(x) \neq 0$ is K and K is compact, there exists an $\epsilon > 0$ such that $|f_2(x)| \geq \epsilon$ for all $x \in K$.

Second, since system 1 is contracting and K is compact, $f_1 \to 0$ uniformly [11]. The transversality condition for the metric M_c is

$$\delta_1' M_1 f_1 + \delta_2' M_2 f_2 = 0 \tag{8}$$

However, since f_2 is bounded below and f_1 converges uniformly to zero, the normal vector to the surface defined by (8) converges to that defined by $\delta_2' M_2 f_2 = 0$ and hence the compact sets of δ of norm one satisfying these conditions converge uniformly.

Now, $d/dt[\delta' M_c \delta] = \delta' H \delta$ where H decomposes into blocks corresponding to δ_1 and δ_2 like so:

$$H = \begin{bmatrix} \alpha(F_1' M_1 + M_1 F_1 + \dot{M}_1) & F_{21}' M_2 + M_1 F_{21} \\ F_{21}' M_1 + M_1 F_{21} & F_{22}' M_2 + M_2 F_{22} + \dot{M}_2 \end{bmatrix}.$$

Consider the fixed $x_1^\star$ to which the contracting system $\dot{x}_1 = f_1(x_1)$ converges, so that $f_1(x^\star = 0)$. Now, since system 1 is contracting, the upper-left block is negative definite and then by the Schur complement it follows that the maximum value of $\delta' H\delta$ on the subspace satisfying $\delta' M f = 0$ can be made strictly negative by choosing α sufficiently large.

Due to continuity of H and the convergence of the sets of δ, this implies that from any initial conditions there exists a finite time after which $\delta' H\delta < 0$ for all δ satisfying (8), which implies the existence of a unique stable periodic orbit by Theorems 1 and 2.
□.

The opposite composition, a transverse-contracting system driving a contracting system clearly converges to a periodic solution due to natural input-to-state stability properties of contracting systems. In a sense, the second system can be considered as being driven by a periodic input.

C. Robustness to Parametric Variation

Suppose the system dynamics depend on some *parameter vector* θ, i.e.

$$\dot{x} = f(x, \theta).$$

When studying robustness of equilibria of such systems, a widely-used method is to search for either a parameter-dependent Lyapunov function via an optimization procedure, see, e.g., [26], [27].

In the context of the present paper, we assume that a particular set K is robustly forward invariant – which can be verified using the methods of [26] – then robust existence of a single globally stable (within K) limit cycle is ensured if one can find a parameter-dependent contraction metric $M(x, \theta)$ which satisfies

$$\delta'(\dot{M}(x,\theta) + 2F(x,\theta)'M(x,\theta) + \lambda M(x,\theta))\delta \leq 0$$

for all δ such that $\delta' M(x,\theta) f(x,\theta) = 0$, and for all $x \in K$ and θ in some set Θ, where K is a forward invariant set.

D. Robustness to Bounded Disturbance

Consider the global coordinates $x_\perp, \tau$ – either implicitly or explicitly defined. Since $\tau \in S^1$ the dynamics of $x_\perp$ can be considered a periodic differential equation with a transformation of time. This makes it clear that any internal perturbation in f which keeps $\dot{\tau} > 0$ and $F_\perp(x)$ contracting still results in a limit cycle (c.f. above).

Bounded external perturbations will also have bounded effect on behavior. Letting $R = \int_{x_1}^{x_2(\cdot)} \delta' M \delta$ we have

$$\dot{R} + 2\lambda R \leq 0$$

Consider a bounded external disturbance, i.e. $\dot{x} = f(x) + d(t)$, where $|d| \leq d_{\max}$, then we have

$$\dot{R} + 2\lambda R \leq |\Pi(x) d(t)|_{M(x)}$$

so after exponentially-forgotten transients, the perturbed system is within a ball of radius R. For further details on such analysis, see [11].

REFERENCES

[1] S. Collins, A. Ruina, R. Tedrake, and M. Wisse, “Efficient bipedal robots based on passive-dynamic walkers,” *Science*, vol. 307, no. 5712, pp. 1082–1085, 2005.

[2] A. Ijspeert, “Central pattern generators for locomotion control in animals and robots: a review,” *Neural Networks*, vol. 21, no. 4, pp. 642–653, 2008.

[3] P. Rapp, “Why are so many biological systems periodic?” *Progress in Neurobiology*, vol. 29, no. 3, pp. 261 – 273, 1987.

[4] J. Mallet-Paret and H. Smith, “The poincaré-bendixson theorem for monotone cyclic feedback systems,” *Journal of Dynamics and Differential Equations*, vol. 2, no. 4, pp. 367–421, 1990.

[5] G. Birkhoff, *Dynamical systems*. Amer. Mathematical Society, 1927.

[6] C. Byrnes, “Topological methods for nonlinear oscillations,” *Notices of the AMS*, vol. 57, no. 9, pp. 1080–1091, 2010.

[7] A. Shiriaev, L. Freidovich, and I. Manchester, “Can we make a robot ballerina perform a pirouette? orbital stabilization of periodic motions of underactuated mechanical systems,” *Annual Reviews in Control*, vol. 32, no. 2, pp. 200 – 211, 2008.

[8] I. R. Manchester, U. Mettin, F. Iida, and R. Tedrake, “Stable dynamic walking over uneven terrain,” *The International Journal of Robotics Research (IJRR)*, vol. 30, no. 3, March 2011.

[9] I. R. Manchester, “Transverse dynamics and regions of stability for nonlinear hybrid limit cycles,” *Proceedings of the 18th IFAC World Congress*, Aug-Sep 2011.

[10] I. R. Manchester, M. Tobenkin, M. Levashov, and R. Tedrake, “Regions of attraction for hybrid limit cycles of walking robots,” in *Proceedings of the IFAC World Congress*, Milan, Italy, 2011.

[11] W. Lohmiller and J. Slotine, “On contraction analysis for non-linear systems,” *Automatica*, vol. 34, no. 6, pp. 683–696, June 1998.

[12] E. Aylward, P. Parrilo, and J. Slotine, “Stability and robustness analysis of nonlinear systems via contraction metrics and SOS programming,” *Automatica*, vol. 44, no. 8, pp. 2163–2170, 2008.

[13] J. K. Hale, *Ordinary Differential Equations*. Robert E. Krieger Publishing Company, New York, 1980.

[14] S. Goodman, “Vector fields with transverse foliations,” *Topology*, vol. 24, no. 3, pp. 333–340, 1985.

[15] X. Yang, “Remarks on three types of asymptotic stability,” *Systems & Control Letters*, vol. 42, no. 4, pp. 299–302, 2001.

[16] G. Leonov, “Generalization of the Andronov-Vitt theorem,” *Regular and Chaotic Dynamics*, vol. 11, no. 2, pp. 281–289, 2006.

[17] I. R. Manchester, M. M. Tobenkin, and J. Wang, “Identification of nonlinear systems with stable oscillations,” in *50th IEEE Conference on Decision and Control (CDC)*. IEEE, 2011.

[18] A. Megretski and A. Rantzer, “System analysis via integral quadratic constraints,” *Automatic Control, IEEE Transactions on*, vol. 42, no. 6, pp. 819–830, 1997.

[19] I. Petersen, V. Ugrinovskii, and A. Savkin, *Robust control design using H^∞ methods*. Springer Verlag, 2000.

[20] M. M. Williamson, “Neural control of rhythmic arm movements,” *Neural Networks*, vol. 11, no. 78, pp. 1379 – 1394, 1998.

[21] W. Wang and J. Slotine, “On partial contraction analysis for coupled nonlinear oscillators,” *Biological cybernetics*, vol. 92, no. 1, pp. 38–53, 2005.

[22] Q.-C. Pham and J.-J. Slotine, “Stable concurrent synchronization in dynamic system networks,” *Neural Networks*, vol. 20, no. 1, pp. 62 – 77, 2007.

[23] M. Morohashi, A. Winn, M. Borisuk, H. Bolouri, J. Doyle, and H. Kitano, “Robustness as a measure of plausibility in models of biochemical networks,” *Journal of theoretical biology*, vol. 216, no. 1, pp. 19–30, 2002.

[24] J. Stelling, U. Sauer, Z. Szallasi, F. Doyle III, and J. Doyle, “Robustness of cellular functions,” *Cell*, vol. 118, no. 6, pp. 675–685, 2004.

[25] U. Jönsson and A. Megretski, “A small-gain theory for limit cycles of systems on lure form,” *SIAM journal on control and optimization*, vol. 44, p. 909, 2005.

[26] W. Tan and A. Packard, “Stability region analysis using polynomial and composite polynomial lyapunov functions and sum-of-squares programming,” *Automatic Control, IEEE Transactions on*, vol. 53, no. 2, pp. 565–571, 2008.

[27] S. Prajna, A. Jadbabaie, and G. Pappas, “A framework for worst-case and stochastic safety verification using barrier certificates,” *Automatic Control, IEEE Transactions on*, vol. 52, no. 8, pp. 1415–1428, 2007.